

Causes

In this issue I will give some of the causes of cracks that I have come across and show some methods which I have used to treat them.

Overgrown feet / Hoof Imbalance



This is by far the most common cause of cracks of the Hoof. During locomotion a balanced Hoof Capsule is essential for shock absorption and fluent blood flow .Should the hoof be excessively out of balance then upon landing on a firm surface abnormal load bearing will transfer excessive force to the **Coronary Band**, causing a disruption to the **keratinisation synthesis** .

It only needs one papillae not to nourish a **Horn Tubule** for a second to start a crack.

Alternatively in overgrown feet a stone trapped between the junction of the sole and the wall will have the effect of driving a wedge into the hoof causing it to split from the bottom.

Trimming Treatment

The treatment in this case is relatively simple, just have the Farrier trim the hoof, relieving the ground pressure from the base of the crack and in most cases where the Coronary Band has not been severed a full recovery is normal if regular trimming is now adhered to.

Conformational Imbalance

Abnormal conformation is a primary cause of hoof cracks. Either Angular Limb deviations or Joint Rotational problems will cause the leg to move forward abnormally .This is commonly known as "Dishing "or "Winging", if we consider the leg as a pendulum ,then the hoof on the end will be moulded to suit the load placed on it from above and will migrate inwards ,outwards away from the direct load. These animals are often referred to as **Pigeon Toed** or **Splay Footed**, when in fact the problem is at the knee or fetlock.

Should the hoof not be trimmed to facilitate a flat, level footfall then even a prolonged trot down the road or a canter in a firm field can lead to a crack developing, usually between the middle of the hoof and the heel as the horse lands on the high side and loads excessively on the other. This is known as a **Quarter Crack**.



Trimming Treatment

In Conformational related cracks, trimming and shoeing to achieve level footfall is fundamental. With a slight **Lateral** (outside first) landing pattern being generally the accepted norm.

Trimming in such a manner often leads to the hoof capsule initially not being trimmed to what is deemed to be aesthetically pleasing to allow for normal landing and loading , and the Farrier should explain this to the owner ,

Abcessation,Treads,Overreaches and other acute Trauma's to the Coronary Band

As explained in previous paragraphs any disruption to the blood supply of the **Coronary Band** will result in a **Hoof Crack**. Should the horse tread on itself, or as was more common in the days of Carriage Horses another horse did the damage. Then a crack will be the result. This will often take the form of a longitudinal split running the length of the wound and whilst being very painful and bloody initially, will normally heal quite well. Veterinary treatment is required for these wounds, to determine that there is no foreign body trapped in the hoof and administer painkillers and often antibiotics

Generally after the initial wound has healed the split will grow down the hoof causing no problem. Exceptions to this are when the crack has grown down and is long enough to be crimped together during load bearing thereby tearing the hoof away from the bone within. With modern hoof glues the Farrier can join top and bottom together allowing normal pain free work.

Should the **Coronary Band** be cut right through (severed), then the condition is called a **False Quarter** regardless of where the wound occurs on the Coronary Band.

The prognosis for this is guarded as the new horn which grows does not have the same **Keratinsation** properties as before and may be prone to **abscess**.

Front and Hind feet shod with Heart Bars. (Horizontal Line above the crack shows that healing has started.

Concave Heartbar My Preferred Shoe for hoof cracks

After the hoof has been trimmed to accommodate the limb conformation. My shoe of choice would be a **Heartbar**. This is a shoe that when properly applied, taking account of the sensitive structures within the hoof, will stabilise movement at the **Coronary Band**.



Hoof nailed and Staped and, trimmed



12 months later!!



NOT Ideal Turnout

It must be stressed however that restricted turnout is advised along with keeping the horse on as flat a surface as possible. If the animal is allowed free range, especially in soft conditions the upward pressure on the sole causes movement at the site of the crack keeping the healing cells apart.

Remember no glue or screws or clips will fix a cracked hoof not correctly trimmed and balanced to the limb.

It is important for a prospective buyer to be aware that although they have the best **Farrier** in the land that can stabilise all cracks, long standing cracks can do irreparable damage to the **Coffin Bone** inside the **Hoof**. This is due to the constant tearing of the **Laminae** from the bone during weightbearing. While they will be fixed externally, dramatic internal bone erosion may have occurred and I would always advise an **x-ray** in such cases.